



ORD

| ORDERED | | Thrombolysis Orders for Dr David Morgan, Dr Mark Ellis & Dr Joseph Kelly |
|---------|------|--|
| Date | Time | |
| | | Admit to: Critical Care Unit |
| | | Diagnosis: |
| | | Condition: |
| | | Allergies: |
| | | Activity: Strict Bedrest |
| | | May elevate HOB maximum of 30° |
| | | Keep affected limbs straight |
| | | May log roll side to side but do not impede blood flow to affected extremity |
| | | Diet: <input type="checkbox"/> Clear Liquids <input type="checkbox"/> Full Liquids <input type="checkbox"/> Regular <input type="checkbox"/> ADA <input type="checkbox"/> NPO |
| | | Lab: Fibrinogen every 6 hr. Notify MD or NP if < 200 |
| | | BMP, CBC, PT/PTT every am |
| | | X-ray: CXR on admission |
| | | Nursing: Vital signs, pulse check and sheath site check every 15 min for 1 hr; then every 1 hr |
| | | Foley catheter pm |
| | | Observe closely for any signs of bleeding |
| | | NO IM injections |
| | | Notify MD or NP of any sudden increase in pain, loss of pulses or mental changes |
| | | Medication: 1) <input type="checkbox"/> Retaplast Recombinate (Retavase) 10 units per 500 ml per hr via infusion catheter |
| | | <input type="checkbox"/> Alteplase (TPA) 10 mg per 500ml NS at _____ ml per hr per sheath |
| | | <input type="checkbox"/> Urokinase 250,000 units in 250 ml NS at 125 ml per hr for 2 hr then 50 ml per hr via infusion catheter |
| | | 2) Heparin 4000 units per 1 liter NS at _____ ml per hr per sheath |
| | | 3) Eptifibatide (Integrilin) bolus & infusion at low dose (1.0 mcg/kg/min) per peripheral line |
| | | 4) Cefazolin (Ancef) 1 gram IVPB every 8 hr |
| | | 5) Meperidine (Demerol) 50 mg IV every 2 hr prn pain |
| | | 6) Promethazine (Phenergan) 12.5 mg IV every 8 hr prn nausea and/or vomiting |
| | | 7) Alprazolam (Xanax) 0.5 mg po every 8 hr pm anxiety or for rest |
| | | 8) Acetaminophen (Tylenol) gr X po every 4 hr pm lesser pain or temperature greater than 100.0 |
| | | 9) Pantopazole Sodium (Protonix) 40 mg po every day |
| | | 10) Identify and list home meds |
| | | 11) Do not use sheaths or infusion catheters for administration of any other meds or to draw blood |
| | | 12) If medications to sheath or infusion catheter is stopped for any reason keep open with NS at 50ml per hr |
| | | MD Signature: |
| | | Date/Time: |



THROMBOLYSIS ORDERS

David Morgan, MD - Mark Ellis, MD - Alice Naquin, CFNP - Joseph Kelly, MD
Form # 18-114.8 (03/09)

▼ Addressograph / Patient Label ▼