ORD

DATE ORDERED		PHYSICIANS ORDERS	POST-OP THORACOTOMY, LOBECTOMY, PNEUMONECTOMY ORDERS Dr. Crocker	SIGN, DATE AND TIME ALL ENTRIES
DATE	TIME			
		1.	Admit to CCU.	
			Diagnosis:	
			I & O's	
			VS Q 15 min til stable then Q 1º per ICU protocol.	
			Labs: CBC, BMP, Mg + ABG and Ionized Calcium now and in a.m.	
			Upright CXR now and in a.m.	
			HOB ↑ 30º legs level	
			O2 @ 3LPM NPR if extubated	
			Respirations: IS Q 1º while awake.	
			EZ PAP Q 4º (7:00 a.m. – 11:00 p.m.)	
			NPO	
			NG to low suction (if patient intubated)	
			Foley to BSD	
			List and resume Pre-op meds per NG tube Clamp NG x 20 minutes and then unclamp.	
			Keep dressing clean, dry, intact	
			Pleural tube to under water seal at 20 cm H2O suction	
			Consult Dr. _____	
		2.	**MEDICATIONS:**	
			Pantoprazole (Protonix) 40 mg daily	
			Mupirocin (Bactroban) 2% nasal ointment to each nostril q 12 hours	
			½ NS + 20 Meq KCL/ liter @ 125cc/hr x 8 hours and then ☐ to 100 cc/hr.	
			If patient a diabetic begin Dr. Crocker's insulin protocol as needed.	
			Cefazolin (Ancef)_____ grams IVPB Q 8º - D/C within 24 hours of pre op dose	
			If allergic to Ancef or Penicillin give Vancomycin x 24 hours – pharmacy to dose.	
			Morphine 2mg IV Q 2º PRN pain.	
			When patient taking PO Δ pain med toOxycodone 5 mg / Acetaminophen 325 mg (Percocet) one tab PO q 3 hours prn pain,	
			may increase to 2 tabs if necessary for pain control.	
			Ondansetron (Zofran)4 mg IV Q 4-6º PRN nausea and vomiting	
			a. ELECTROLYTE REPLACEMENT THERAPY:	
			1. Supplemental Potassium Chloride:	
			a. If Potassium level less than 3.5 give 60 mEq IVPB, infuse over 1 ½ hours	
			b. If Potassium level is between 3.5 and 4.0 give 40 mEq IVPB, infuse over 1 hour.	
			c. Repeat Potassium level in 1 hour post infusion.	
			2. Supplemental Magnesium:	
			a. Mix 2 grams Magnesium Sulfate in 100 ml D5W	
			1. If Magnesium level is less than or equal 2 mg per dl give 2 grams over 30 minutes	
			2. Repeat Magnesium level in 1 hour post infusion	

Physician Signature_____ Date/Time _____

BAPTIST MEMORIAL HOSPITAL GOLDEN TRIANGLE

POST-OP THOROCOTOMY, LOBECTOMY, PNEUMONECTOMY ORDERS - DR. CROCKER

▼ Addressograph / Patient Label ▼

ORD

DATE ORDERED		PHYSICIANS ORDERS	POST-OP THORACOTOMY, LOBECTOMY, PNEUMONECTOMY ORDERS Dr. Crocker	SIGN, DATE AND TIME ALL ENTRIES
DATE	TIME			
		3. Supplemental Calcium Chloride:		
		a. If Ionized Calcium level less than 4.0 mg/dl give 1 gram Calcium Chloride in 50 ml D5W over 1 hour		
		b. Repeat Calcium level in 1 hour post infusion		
		3. HEMODYNAMIC:		
		a. VASODILATORS:		
		_____ If Systolic BP greater than 160 and SVR less than 1500 dynes begin Nitroglycerin drip 50 mg in 250cc of D5W initiate at 5 micrograms per min and titrate not to exceed 100 micrograms per min (30 ml per hr)		
		_____ If Systolic BP remains greater than 160 with titrated Nitroglycerine drip at 5 micrograms per kilogram per minute (100 micrograms per minute). If Systolic BP remains greater than 160 with titrated Nitroglycerin drip at 5 Micrograms per kilogram per minute (100 micrograms per minute) start Nitroprusside (Nipride) drip 100 mg in 250 ml D5W initiate at 0.1 – 0.3 micrograms per kilogram per minute not to exceed 10 micrograms per kilogram per minute.		
		Other Drips:_____		
		b. VOLUME REPLACEMENT: (notify surgeon with the intervention)		
		1. If Systolic BP less than 90 and the PAD less than 12 or CVP less than 6		
		a._____ Infuse Hespan 500 ml bolus		

Physician Signature_____ Date/Time _____

BAPTIST MEMORIAL HOSPITAL GOLDEN TRIANGLE

POST-OP THOROCOTOMY, LOBECTOMY, PNEUMONECTOMY ORDERS - DR. CROCKER

▼ Addressograph / Patient Label ▼