

UROLOGY OUTPATIENT ADMIT ORDERS



Check appropriate boxes:

Primary diagnosis: _____

Operative Permit for: _____

DIET: NPO

Vital Signs: routine

INT IV fluid: _____ Rate: _____ ml/hr

Pre-op studies:

- CBC with diff CBC CMP BMP UA PT/PTT
 Sterile midstream urine collection for culture and sensitivity
 EKG CXR: PA and lateral Type and Screen Other: _____

Beta Blocker Pre OP

Upon admission to outpatient surgery (via outpatient surgery), outpatient RN to document last dose taken. If >24 hours since last dose, prior to anticipated surgery time, give _____ with sips of water.

Hold beta blocker for pulse <50 and/or systolic BP <100.

PreOP Medications: GIVE WITHIN ONE HOUR OF INCISION

Transrectal Prostate Biopsy:

- Levofloxacin 500 mg PO
 Levofloxacin 500 mg IV
 For patient <80 kg: Cefoxitin (Mefoxin) 1 gram IV
 For patient >80 kg: Cefoxitin (Mefoxin) 2 gram IV
 Beta-lactam (PCN or cephalosporin) allergy:
 Gentamicin _____(rec. 1mg/kg) IV+ Metronidazole 500 mg IV
 Gentamicin _____(rec. 1 mg/kg) IV + Clindamycin 600 mg IV

Penile Prosthesis (insertion, removal or revision):

- Ampicillin/sulbactam (Unasyn) 3 gm IV
 For patient < 80 kg: Cefazolin (Ancef) 1 gram IV + Gentamicin _____(rec 1mg/kg)
 For patient > 80 kg: Cefazolin (Ancef) 2 gram IV + Gentamicin _____(rec 1mg/kg)
 Beta-lactam (PCN or cephalosporin) allergy:
 Gentamicin _____(rec 1 mg/kg) IV+ Vancomycin _____(rec 15 mg/kg)IV
 Gentamicin _____(rec 1 mg/kg) IV + Clindamycin 600 mg IV

Pubovaginal Sling

- For patient < 80 kg: Cefazolin (Ancef) 1 gram IV
 For patient > 80 kg: Cefazolin (Ancef) 2 gram IV
 Levofloxacin 500 mg IV
 Other: _____
 Beta-lactam (PCN or cephalosporin) allergy:
 Gentamicin _____(rec. 1mg/kg) IV+ Metronidazole 500 mg IV
 Gentamicin _____(rec. 1 mg/kg) IV + Clindamycin 600 mg IV

Physician Signature: _____ Date/Time _____



▼ Patient Label ▼

UROLOGY OUTPATIENT ADMIT ORDERS



UROLOGY OUTPATIENT ADMIT ORDERS

Cystoscopy alone (urine culture positive or unavailable, pre-operative catheter, placement of prosthetic material or Cystoscopy with Manipulation of Upper Urinary Tract, Shockwave Lithotripsy, Ureteroscopy)

- Levofloxacin 500mg PO
- Levofloxacin 500mg IV
- Trimethoprim – Sulfamethoxazole DS one tablet PO
- Other: _____

TURP (positive urine culture, no culture obtained, or catheter in place):

- For patient <80 kg: Cefazolin (Ancef) 1 gram IV + Gentamicin _____(rec 1 mg/kg) IV
- For patient >80 kg: Cefazolin (Ancef) 2 gram IV + Gentamicin _____(rec 1 mg/kg) IV
- Levofloxacin 500mg PO
- Levofloxacin 500mg IV

Preop Antibiotics for Other Procedures: _____

Other Orders: _____

Physician Signature: _____ Date/Time _____



▼ Patient Label ▼

UROLOGY OUTPATIENT ADMIT ORDERS